



Decision Support

# An axiomatization of entropy of capacities on set systems

Aoi Honda <sup>a,b,\*</sup>, Michel Grabisch <sup>b</sup>

<sup>a</sup> *Kyushu Institute of Technology, Japan*

<sup>b</sup> *Université de Paris I – Panthéon-Sorbonne, France*

Received 7 October 2005; accepted 19 June 2007

Available online 29 June 2007

---

## Abstract

We present an axiomatization of the entropy of capacities defined on set systems which are not necessarily the whole power set, but satisfy a condition of regularity. This entropy encompasses the definition of Marichal and Roubens for the entropy of capacities. Its axiomatization is in the spirit of the one of Faddeev for the classical Shannon entropy. In addition, we present also an axiomatization of the entropy for capacities proposed by Dukhovny.  
© 2007 Elsevier B.V. All rights reserved.

*Keywords:* Entropy; Capacity; Regular set system; Generalized Faddeev's axiom

---

## 1. Introduction

The notion of entropy of a probability measure, as proposed by Shannon [11], is of primary importance in many application fields, e.g., information theory, operations research, classification, data mining, etc.

It was therefore natural to investigate about the entropy of capacities [1] (also called fuzzy measures [12], non-additive measures [2]), since they are a generalization of probability measures. After a first attempt by Yager [13], Marichal and Roubens [10,9] proposed a definition having suitable properties, and which can be considered as the generalization of the Shannon entropy. Another attempt was also done by Dukhovny [3] in a different spirit. All these works considered finite universal sets, and the power set as underlying set system. In a previous paper [6], we proposed a definition of entropy to be able to work with capacities defined on set systems which are not necessarily the whole power set, but satisfy a regularity condition. Our definition encompasses one of the Marichal and Roubens, and also encompasses the Shannon entropy. Our entropy can be applied to capacities on lattices as well.

An important question about entropy is to find suitable axiomatizations, where axioms are enough understandable or natural. There exist many of them for the classical Shannon entropy (see, e.g., [4]). For capacities,

---

\* Corresponding author. Address: Kyushu Institute of Technology, 680-4 Kawazu, Iizuka 820-8502, Japan. Tel.: +81 948 29 7730; fax: +81 948 29 7709.

E-mail address: [aoi@ces.kyutech.ac.jp](mailto:aoi@ces.kyutech.ac.jp) (A. Honda).

to the best of our knowledge, there exists only the work of Kojadinovic et al. [7]. Their axiomatization is however rather complicated, due to the presence of a recursive axiom, whose meaning is hard to grasp.

Our aim in this paper is to present a new axiomatization which is more understandable in our general framework where the set system is not necessarily the whole power set, but a subset of it. Our axiomatization is in the spirit of the one proposed by Faddeev [5] for the Shannon entropy.

The paper is organized as follows. Section 2 introduces notations and necessary materials about set systems. Then Section 3 presents capacities on set systems and the definition of their entropy. The axiomatization of entropy is given in Section 4, and proofs of the main theorems are given in Section 5. Lastly, we give some final remarks on the suitability of regular set systems as a fundamental concept, and some directions for future research in Section 6.

## 2. Mathematical background

We begin by introducing some notations and definitions.

Throughout this paper, we consider a finite universal set  $N := \{1, 2, \dots, n\}$ ,  $n \geq 2$ , and  $2^N$  denotes the power set of  $N$ . Let us consider  $\mathcal{N}$  as a subcollection of  $2^N$  which contains  $\emptyset$  and  $N$ . Then we call  $(N, \mathcal{N})$  (or simply  $\mathcal{N}$  if no confusion occurs) a *set system*.

A set system endowed with inclusion is a particular case of a *partially ordered set*  $(P, \leq)$ , i.e., a set  $P$  endowed with a partial order (reflexive, antisymmetric and transitive)  $\leq$ , as for example  $(\mathcal{N}, \subseteq)$ .

Let  $A, B \in \mathcal{N}$ . We say that  $A$  is *covered* by  $B$ , and write  $A \prec B$  or  $B \succ A$ , if  $A \subsetneq B$  and  $A \subseteq C \subsetneq B$  together with  $C \in \mathcal{N}$  imply  $C = A$ .

**Definition 1 (Maximal chain of set system).** Let  $\mathcal{N}$  be a set system. We call  $C = (c_0, c_1, \dots, c_m)$  a *maximal chain* of  $\mathcal{N}$  if  $C$  satisfies  $\emptyset = c_0 \prec c_1 \prec \dots \prec c_m = N$ ,  $c_i \in \mathcal{N}$ ,  $i = 0, \dots, m$ .

We denote the set of all maximal chains of  $\mathcal{N}$  by  $\mathcal{C}(\mathcal{N})$ .

**Definition 2 (Totally ordered set system).** We say that  $(N, \mathcal{N})$  is a *totally ordered set system* if for any  $A, B \in \mathcal{N}$ , either  $A \subseteq B$  or  $A \supseteq B$ .

If  $(N, \mathcal{N})$  is a totally ordered set system, then  $|\mathcal{C}(\mathcal{N})| = 1$ .

**Definition 3 (Regular set system [6]).** We say that  $(N, \mathcal{N})$  is a *regular set system* if for any  $C \in \mathcal{C}(\mathcal{N})$ , the length of  $C$  is  $n$ , i.e.  $|C| = n + 1$ .

Equivalently,  $\mathcal{N}$  is a regular set system if and only if  $|A \setminus B| = 1$  for any  $A, B \in \mathcal{N}$  satisfying  $A \succ B$ .

## 3. Entropy of a capacity

**Definition 4 (Capacity on a set system).** Let  $(N, \mathcal{N})$  be a set system. A function  $v : \mathcal{N} \rightarrow [0, 1]$  is a *capacity* on  $(N, \mathcal{N})$  if it satisfies  $v(\emptyset) = 0$ ,  $v(N) = 1$  and, for any  $A, B \in \mathcal{N}$ ,  $v(A) \leq v(B)$  whenever  $A \subseteq B$ .

Let  $v$  be a capacity on  $(N, \mathcal{N})$ . For  $C := (c_0, c_1, \dots, c_m) \in \mathcal{C}(\mathcal{N})$ , define  $p^{v,C}$  by

$$p^{v,C} := (p_1^{v,C}, p_2^{v,C}, \dots, p_m^{v,C}) = (v(c_1) - v(c_0), v(c_2) - v(c_1), \dots, v(c_m) - v(c_{m-1})).$$

Note that  $p^{v,C}$  satisfies  $p_i^{v,C} \geq 0$ ,  $i = 1, \dots, m$ , and  $\sum_{i=1}^m p_i^{v,C} = 1$ .

We introduce further concepts about capacities, which will be useful for stating axioms.

**Definition 5 (Dual capacity).** Let  $v$  be a capacity on  $(N, \mathcal{N})$ . Then the *dual capacity* of  $v$  is defined on  $\mathcal{N}^d := \{A^c \in 2^N \mid A \in \mathcal{N}\}$  by  $v^d(A) := 1 - v(A^c)$  for any  $A \in \mathcal{N}^d$ , where  $A^c := N \setminus A$ .

A Hasse diagram for  $\mathcal{N}^d$  is obtained by turning upside down a Hasse diagram for  $\mathcal{N}$  (cf. Fig. 1).

**Definition 6 (Permutation of  $v$ ).** Let  $\pi$  be a permutation on  $N$ . Then the *permutation* of  $v$  by  $\pi$  is defined on  $\pi(\mathcal{N}) := \{\pi(A) \in 2^N \mid A \in \mathcal{N}\}$  by  $\pi \circ v(A) := v(\pi^{-1}(A))$ .




Fig. 1. Dual set system.




Fig. 2. Permutation of set system.

**Example 7.** We show an example of a permutation of a set system. Let  $N := \{1, 2, 3\}$  and  $\mathcal{N} := \{\emptyset, \{1\}, \{3\}, \{1, 2\}, \{1, 3\}, \{2, 3\}, N\}$  and let  $\pi = \begin{pmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \end{pmatrix}$ . Then, for instance  $\pi \circ v(\{2, 3\}) := v(\pi^{-1}(\{2, 3\})) = v(\{1, 2\})$  (cf. Fig. 2). For the maximal chain  $C := (\emptyset, \{2\}, \{2, 3\}, N) \in \mathcal{C}(\pi(\mathcal{N}))$ ,  $\pi^{-1}(C) = (\pi^{-1}(\emptyset), \pi^{-1}(\{2\}), \pi^{-1}(\{2, 3\}), \pi^{-1}(N)) = (\emptyset, \{1\}, \{1, 2\}, N) \in \mathcal{C}(\mathcal{N})$ .

As it is usual for functions, we denote the *restriction* of  $v$  to some set system  $\mathcal{N}' \subseteq \mathcal{N}$  by  $v|_{\mathcal{N}'}$ , i.e.,  $v|_{\mathcal{N}'}(A) := v(A)$  for any  $A \in \mathcal{N}'$ .

Let us consider a chain of length 2 as a set system, denoted by  $\mathbf{2}$  (e.g.  $\{\emptyset, \{1\}, \{1, 2\}\}$ ), and a capacity  $v^2$  on it. We denote by the triplet  $(0, u, 1)$ ,  $0 \leq u \leq 1$ , the values of  $v^2$  along the chain.

**Definition 8 (Embedding of  $v^2$ ).** Let  $v$  be a capacity on a totally ordered regular set system  $(N, \mathcal{N})$ , where  $\mathcal{N} := \{c_0, \dots, c_n\}$  such that  $c_{i-1} \prec c_i$ ,  $i = 1, \dots, n$ , and let  $v^2 := (0, u, 1)$  be a capacity on  $\mathbf{2}$ . Then for  $c_k \in \mathcal{N}$ ,  $v^{c_k}$  is called the *embedding* of  $v^2$  into  $v$  at  $c_k$ , and defined on the totally ordered regular set system  $(N^{c_k}, \mathcal{N}^{c_k})$  by

$$v^{c_k}(A) := \begin{cases} v(c_{k-1}) + u \cdot (v(c_k) - v(c_{k-1})), & \text{if } A = c'_k, \\ v(c_j), & \text{if } A = c_j \text{ for } j < k, \text{ or } A = c''_j \text{ for } j \geq k \end{cases} \quad (3.1)$$

with  $\{i_k\} := c_k \setminus c_{k-1}$ ,  $N^{c_k} := \{1, \dots, i_{k-1}, i'_k, i''_k, i_{k+1}, \dots, n\}$  with  $i'_k \neq i''_k$ ,  $(N \setminus \{i_k\}) \cap \{i'_k, i''_k\} = \emptyset$ ,  $c'_k := (c_k \setminus \{i_k\}) \cup i'_k$ ,  $c''_j := (c_j \setminus \{i_k\}) \cup \{i'_k, i''_k\}$  for  $j = k, \dots, n$ , and  $\mathcal{N}^{c_k} := \{c_0, \dots, c_{k-1}, c'_k, c''_k, c''_{k+1}, \dots, c''_n\}$ .

**Example 9.** Let us take  $N := \{1, 2, 3\}$ ,  $\mathcal{N} := \{\emptyset, 1, 12, 123\}$  and  $c_k = 12$  (“12” stands for  $\{1, 2\}$ , etc.). Then  $i_k = 2$ , which is replaced by  $2'$  and  $2''$ . We obtain

$$\begin{aligned} N^{12} &= \{1, 2', 2'', 3\}, \\ \mathcal{N}^{12} &= \{\emptyset, 1, 12', 12'2'', 12'2''3\}, \\ v^{12}(1) &= v(1), \end{aligned}$$

$$v^{12}(12') = v(1) + u \cdot (v(12) - v(1)),$$

$$v^{12}(12'2'') = v(12),$$

$$v^{12}(12'2''3) = v(123).$$

We now turn to the definition of the entropy. We first recall the classical definition of Shannon.

**Definition 10 (Shannon entropy).** Let  $p = (p_1, \dots, p_n)$  be a probability measure on  $N$ . Then the *Shannon entropy* of  $p$  is defined by

$$(S) \quad H_S(p) = H_S(p_1, \dots, p_n) := - \sum_{i=1}^n p_i \log p_i,$$

where  $\log$  denotes the base 2 logarithm, and by convention  $0 \log 0 := 0$ .

**Definition 11 (Entropy of a capacity on a set system [6]).** Let  $v$  be a capacity on a regular set system  $(N, \mathcal{N})$ . The entropy of  $v$  is defined by

$$(HG) \quad H_{HG}(v) := \frac{1}{|\mathcal{C}(\mathcal{N})|} \sum_{C \in \mathcal{C}(\mathcal{N})} H_S(p^{v,C}).$$

We recall the definition of Dukhovny [3] for capacities defined on the power set, which we extend to our framework. We call it minimum entropy.

**Definition 12.** Let  $v$  be a capacity on a regular set system  $(N, \mathcal{N})$ . The *minimum entropy* of  $v$  is defined by

$$(D) \quad H_D(v) := \min_{C \in \mathcal{C}(\mathcal{N})} H_S(p^{v,C}).$$

As it is clear from its definition,  $H_D$  is not differentiable.

Before entering the axiomatization part, a remark on the domain of entropy functions is in order. For mathematical rigor, one should define the Shannon entropy relative to the cardinality of the universe, hence when necessary we will use the notation  $H_S^n$ , with  $|N| = n$ . More precisely, let  $\Delta_n$  be the set of all probability measures on  $N = \{1, \dots, n\}$  and algebra  $2^N$ .  $H_S$  is a function defined on  $\Delta := \bigcup_{n=2}^{\infty} \Delta_n$  to  $[0, \infty)$ , and  $H_S^n$  is its restriction to  $\Delta_n$ .

Accordingly, we do the same for our definition, up to the fact that  $H_{HG}$  depends not only on  $N$  but also on the set system  $\mathcal{N}$ . As it will become clear hereafter, in fact only the number of maximal chains matters. But since in the axiomatization part, this cannot be known in advance, we let as superindex the set system  $\mathcal{N}$ , and also the cardinality  $n$  of the universe, the latter only for clarity since this can be known from the set system:  $H_{HG}^{n,\mathcal{N}}$  or  $H^{n,\mathcal{N}}$ . We denote by  $\mathcal{A}_{n,\mathcal{N}}$  the set of all capacities defined on  $(N, \mathcal{N})$ , and the domain of  $H_{HG}$  is  $\mathcal{A} := \bigcup_{n=2}^{\infty} \bigcup_{\mathcal{N} \text{ on } N} \mathcal{A}_{n,\mathcal{N}}$ .

#### 4. Axiomatization of the entropy of capacities

First, we recall Faddeev's axiomatization, which will serve as a basis for our axiomatization. In what follows,  $H^n$  is a function from  $\Delta_n$  to  $[0, \infty)$ .

(F1)  $f(x) := H^2(p, 1 - p)$  is continuous on  $0 \leq p \leq 1$ , and there exists  $p_0 \in [0, 1]$  such that  $f(p_0) > 0$ .

(F2) For any permutation  $\pi$  on  $N$ ,

$$H^n(p_{\pi(1)}, \dots, p_{\pi(n)}) = H^n(p_1, \dots, p_n).$$

(F3) If  $p_n = q + r$ ,  $q > 0$ ,  $r > 0$ , then

$$H^{n+1}(p_1, \dots, p_{n-1}, q, r) = H^n(p_1, \dots, p_n) + p_n H^2(q/p_n, r/p_n).$$

**Theorem 13** (Faddeev’s axiomatization of the Shannon entropy [5]). *Under the condition  $H^2(1/2, 1/2) = 1$ ,  $H^n = H_S^n$  for all  $n \geq 2$  holds if and only if (F1), (F2) and (F3) hold.*

We introduce five axioms for the entropy of capacities. In what follows,  $H^{n,\nu}$  is a function from  $\mathcal{A}_{n,\nu}$  to  $[0, \infty)$ .

**(HG1) (Continuity).** The function  $f(u) := H^{2,2}(0, u, 1)$  is continuous on  $[0, 1]$ , and there exists  $u_0 \in [0, 1]$  such that  $f(u_0) > 0$ .

**(HG2) (Dual invariance).** For any capacity  $(0, u, 1)$  on **2**,

$$H^{2,2}(0, u, 1) = H^{2,2}(0, 1 - u, 1).$$

**(HG3) (Increase by embedding).** Let  $v$  be a capacity on a totally ordered regular set system  $(N, \mathcal{N})$  with  $n \geq 2$ . Then for any  $c_k \in \mathcal{N}$  and any  $v^2 := (0, u, 1)$ , the entropy of  $v^{c_k}$  is

$$H^{n+1,\nu^{c_k}}(v^{c_k}) = H^{n,\nu}(v) + (v(c_k) - v(c_{k-1})) \cdot H^{2,2}(0, u, 1). \tag{4.1}$$

**(HG4) (Convexity).** Let  $(N, \mathcal{N})$  with  $n \geq 2$ ,  $(N, \mathcal{N}_1), \dots, (N, \mathcal{N}_k)$  be regular set systems on  $\mathcal{N}$ , satisfying  $\mathcal{C}(\mathcal{N}) = \mathcal{C}(\mathcal{N}_1) \cup \dots \cup \mathcal{C}(\mathcal{N}_k)$ , and  $\mathcal{C}(\mathcal{N}_i) \cap \mathcal{C}(\mathcal{N}_j) = \emptyset$  for all  $i \neq j$ . Then there exist unique  $\alpha_1, \dots, \alpha_k \in ]0, 1[$  satisfying  $\sum_{i=1}^k \alpha_i = 1$  such that for any capacity  $v$  on  $(N, \mathcal{N})$ ,

$$H^{n,\nu}(v) = \alpha_1 H^{n,\nu_1}(v|_{\mathcal{N}_1}) + \dots + \alpha_k H^{n,\nu_k}(v|_{\mathcal{N}_k}).$$

**(HG5) (Permutation invariance).** Let  $v$  be a capacity on a regular set system  $\mathcal{N}$  with  $n \geq 2$ . Then for any permutation  $\pi$  on  $N$  satisfying  $\pi(\mathcal{N}) = \mathcal{N}$ , it holds that

$$H^{n,\nu}(v) = H^{n,\nu}(\pi \circ v).$$

**Theorem 14.** *Under the condition  $H^{2,2}(0, 1/2, 1) = 1$ ,  $H^{n,\nu} = H_{HG}^{n,\nu}$  for  $n \geq 2$  and any regular set system  $\mathcal{N}$  on  $N$ , if and only if (HG1), (HG2), (HG3), (HG4) and (HG5) hold.*

(See proof in Section 5)

For additive capacities on  $2^N$ ,  $H_{HG}$  is equal to the Shannon entropy  $H_S$ . We discuss in detail the above axioms, in the light of Faddeev’s axioms.

- **Continuity:**

We have  $f(u) = H_{HG}^{2,2}(0, u, 1) = H_S^2(p^{(0,u,1),C}) = H_S^2(u, 1 - u)$ , where  $C := (\emptyset, \{1\}, \{1, 2\})$ . Therefore, (HG1) corresponds to (F1).

- **Dual invariance:**

More generally, even where  $v$  is defined on a regular set system which is not necessarily **2**,  $H_{HG}(v)$  is dual invariant.

**Proposition 15 (Dual invariance).** *For any capacity  $v$  on a regular set system,  $H_{HG}(v^d) = H_{HG}(v)$  holds.*

**Proof.** Let  $v$  be a capacity on  $\mathcal{N}$ . For any  $a \in \mathcal{N}$ ,  $(a^c)^c = a$ , hence the dual mapping is a bijection from  $\mathcal{N}$  to  $\mathcal{N}^d$ . Then,  $C := (c_0, \dots, c_n) \in \mathcal{C}(\mathcal{N})$  if and only if  $C^d := (c_n^c, \dots, c_0^c) \in \mathcal{C}(\mathcal{N}^d)$ , since  $c_i < c_{i+1}$  implies  $c_i^c > c_{i+1}^c$ . Hence  $|\mathcal{C}(\mathcal{N})| = |\mathcal{C}(\mathcal{N}^d)|$ . In addition, we have for any  $C \in \mathcal{C}(\mathcal{N})$ ,

$$\begin{aligned} H_S(p^{v,C}) &= \sum_{i=1}^n h[v(c_i) - v(c_{i-1})] = \sum_{i=1}^n h[(1 - v(c_{i-1})) - (1 - v(c_i))] = \sum_{i=1}^n h[v^d(c_{i-1}^c) - v^d(c_i^c)] \\ &= H_S(p^{v^d,C^d}), \end{aligned}$$

where  $h(x) := -x \log x$ . Therefore,

$$H_{HG}(v^d) = \frac{1}{|\mathcal{C}(\mathcal{N}^d)|} \sum_{C \in \mathcal{C}(\mathcal{N}^d)} H_S(p^{v^d,C^d}) = \frac{1}{|\mathcal{C}(\mathcal{N})|} \sum_{C^d \in \mathcal{C}(\mathcal{N}^d)} H_S(p^{v,C}) = H_{HG}(v). \quad \square$$

The Shannon entropy of a probability measure satisfies dual invariance, since a probability measure and its dual measure are identical.

• *Increase by embedding:*

Let  $v$  be a capacity on a totally ordered regular set system  $\mathcal{N} = \{C := \{c_0, c_1, \dots, c_n\}\}$ , where  $c_{i-1} \prec c_i$ ,  $i = 0, \dots, n$ , and consider the embedding of  $v^2 := (0, u, 1)$  into  $v$  at  $c_k$ . Then

$$H_{HG}(v^{c_k}) = H_S(p^{v^{c_k}, C'}),$$

where  $C' := (c_0, \dots, c_{k-1}, c'_k, c''_k, c''_{k+1}, \dots, c''_n)$ , with  $c'_k := (c_k \setminus \{i_k\}) \cup i'_k$ ,  $c''_j := (c_j \setminus \{i_k\}) \cup \{i'_k, i''_k\}$ ,  $j = k, \dots, n$  (see Definition 8). And by (F3), we have

$$H_S(p^{v^{c_k}, C'}) = H_S(p^{v, C}) + (v(c_k) - v(c_{k-1})) \cdot H_S(u, 1 - u)$$

which can be rewritten as

$$H_{HG}(v^{c_k}) = H_{HG}(v) + (v(c_k) - v(c_{k-1})) \cdot H_{HG}(v^2).$$

This is exactly (HG3).

The embedding of  $(0, u, 1)$  at  $c_k \in \mathcal{N}$  means that  $i_k := c_k \setminus c_{k-1} \in N$  is splitted into  $i'_k, i''_k$ . In other words,  $i_k$  is the union of  $i'_k$  and  $i''_k$ . Hence (HG3) means that for capacities on totally ordered regular set systems, if elements of  $N$  are splitted, then the entropy increases according to (4.1).

• *Convexity:*

This axiom says that the computation of the entropy follows the structure of the set system, whatever the capacity is. If the (regular) set system can be decomposed into several (regular) subsystems, provided all elements of  $N$  are present in all subsystems, and that no redundancy occurs (no common maximal chain), then the computation can also be decomposing into these subsystems, each corresponding to the entropy computed on a subsystem. It is important to note that the way of decomposing the computation does not depend on the capacity but only on the decomposing, i.e., the  $\alpha_i$ 's are independent of the capacity.

• *Permutation invariance:*

(HG5) corresponds to (F2). Note that there are set systems such that no non-identity permutation makes them invariant.

To finish this section, we consider a modification of our axiomatization so as to recover the entropy defined by Dukhovny [3]. We modify (HG4) as follows:

(HG4') Let  $(N, \mathcal{N}), (N, \mathcal{N}_1), \dots, (N, \mathcal{N}_k)$  be regular set systems satisfying  $\mathcal{C}(\mathcal{N}) = \mathcal{C}(\mathcal{N}_1) \cup \dots \cup \mathcal{C}(\mathcal{N}_k)$ . Then for any capacity  $v$  on  $\mathcal{N}$ ,

$$H^{n,v}(v) = \min \left\{ H^{n,v_1}(v|_{\mathcal{N}_1}), \dots, H^{n,v_k}(v|_{\mathcal{N}_k}) \right\}.$$

**Theorem 16.** Under the condition  $H^{2,2}(0, 1/2, 1) = 1$ ,  $H^{n,\mathcal{N}} = H_D^{n,\mathcal{N}}$  for all  $n \geq 2$  and all regular set systems  $\mathcal{N}$  on  $N$  holds if and only if (HG1), (HG2), (HG3) and (HG4') hold.

(See proof in Section 5)

**5. Proof of Theorems 14 and 16**

**Proof of Theorem 14 (Necessity).** We have  $H_{HG}^{2,2}(0, u, 1) = -u \log u - (1 - u) \log(1 - u)$ , hence obviously  $H_{HG}$  satisfies (HG1).

By Proposition 15,  $H_{HG}$  satisfies (HG2).

We show that  $H_{HG}$  satisfies (HG3). Let  $v$  be a capacity on  $\mathcal{N} := \{C := (c_0, \dots, c_n)\}$  such that  $c_{i-1} \prec c_i$ ,  $i = 1, \dots, n$ . Then the capacity  $v^{c_k}$  which is embedding of  $(0, u, 1)$  into  $v$  at  $c_k$  is defined in  $\mathcal{N}^{c_k} = \{C' := \{c_0 \prec \dots \prec c_{k-1} \prec c'_k \prec c''_k \prec \dots \prec c''_n\}\}$  with notations of Definition 8. Then we have

$$\begin{aligned} H_{HG}^{n+1, v^{c_k}}(v^{c_k}) &= H_S(p^{v^{c_k}, C'}) = \sum_{i=1}^{k-1} h[v^{c_k}(c_i) - v^{c_k}(c_{i-1})] + h[v^{c_k}(c'_k) - v^{c_k}(c_{k-1})] + h[v^{c_k}(c''_k) - v^{c_k}(c'_k)] \\ &\quad + \sum_{i=k+1}^n h[v^{c_k}(c''_i) - v^{c_k}(c''_{i-1})] \\ &= \sum_{i \neq k} h[v(c_i) - v(c_{i-1})] + (v(c_k) - v(c_{k-1})) \left( h \left[ \frac{v^{c_k}(c'_k) - v^{c_k}(c_{k-1})}{v(c_k) - v(c_{k-1})} \right] + h \left[ \frac{v^{c_k}(c''_k) - v^{c_k}(c'_k)}{v(c_k) - v(c_{k-1})} \right] \right) \\ &\quad + h[v^{c_k}(c'_k) - v^{c_k}(c_{k-1})] \\ &= \sum_{i=1}^n h[v(c_i) - v(c_{i-1})] + (v(c_k) - v(c_{k-1})) H_{HG}^{2,2} \left( 0, \frac{v^{c_k}(c'_k) - v^{c_k}(c_{k-1})}{v(c_k) - v(c_{k-1})}, 1 \right) \\ &= H_{HG}^{n, v}(v) + (v(c_k) - v(c_{k-1})) H_{HG}^{2,2}(0, u, 1), \end{aligned}$$

where  $u = (v^{c_k}(c'_k) - v^{c_k}(c_{k-1})) / (v(c_k) - v(c_{k-1}))$ .

We show that  $H_{HG}$  satisfies (HG4). We have, for set systems  $\mathcal{N}_1, \mathcal{N}_2, \dots, \mathcal{N}_k$  satisfying conditions of (HG4):

$$\begin{aligned} H_{HG}^{n, v}(v) &= \frac{1}{|\mathcal{C}(\mathcal{N})|} \sum_{C \in \mathcal{C}(\mathcal{N})} H_S^n(p^{v, C}) = \frac{1}{|\mathcal{C}(\mathcal{N})|} \left( \sum_{C \in \mathcal{C}(\mathcal{N}_1)} H_S^n(p^{v, C}) + \dots + \sum_{C \in \mathcal{C}(\mathcal{N}_k)} H_S^n(p^{v, C}) \right) \\ &= \frac{|\mathcal{C}(\mathcal{N}_1)|}{|\mathcal{C}(\mathcal{N})|} \frac{1}{|\mathcal{C}(\mathcal{N}_1)|} \sum_{C \in \mathcal{C}(\mathcal{N}_1)} H_S^n(p^{v|_{\mathcal{N}_1}, C}) + \dots + \frac{|\mathcal{C}(\mathcal{N}_k)|}{|\mathcal{C}(\mathcal{N})|} \frac{1}{|\mathcal{C}(\mathcal{N}_k)|} \sum_{C \in \mathcal{C}(\mathcal{N}_k)} H_S^n(p^{v|_{\mathcal{N}_k}, C}) \\ &= \frac{|\mathcal{C}(\mathcal{N}_1)|}{|\mathcal{C}(\mathcal{N})|} H_{HG}^{n, v|_{\mathcal{N}_1}}(v|_{\mathcal{N}_1}) + \dots + \frac{|\mathcal{C}(\mathcal{N}_k)|}{|\mathcal{C}(\mathcal{N})|} H_{HG}^{n, v|_{\mathcal{N}_k}}(v|_{\mathcal{N}_k}) \end{aligned}$$

and

$$\frac{|\mathcal{C}(\mathcal{N}_1)|}{|\mathcal{C}(\mathcal{N})|} + \dots + \frac{|\mathcal{C}(\mathcal{N}_k)|}{|\mathcal{C}(\mathcal{N})|} = 1.$$

We show that  $H_{HG}$  satisfies (HG5). For any  $C = (c_0, c_1, \dots, c_m) \in \mathcal{C}(\pi(\mathcal{N}))$ ,  $\pi^{-1}(C) := (\pi^{-1}(c_0), \pi^{-1}(c_1), \dots, \pi^{-1}(c_m)) \in \mathcal{C}(\mathcal{N})$ , because for any  $A, B \in \pi(\mathcal{N})$ ,  $A \prec B$  implies  $\pi^{-1}(A) \prec \pi^{-1}(B)$ , and for any  $C_i, C_j \in \mathcal{C}(\mathcal{N})$ ,  $i \neq j$ ,  $\pi(C_i) \neq \pi(C_j)$ . Consequently, we have

$$\begin{aligned} H_{HG}^{n, \pi(\cdot)}(\pi \circ v) &= \frac{1}{|\mathcal{C}(\pi(\mathcal{N}))|} \sum_{C \in \mathcal{C}(\pi(\mathcal{N}))} H_S^n(p^{\pi \circ v, C}) = \frac{1}{|\mathcal{C}(\mathcal{N})|} \sum_{C \in \mathcal{C}(\mathcal{N})} H_S^n(p^{\pi \circ v, \pi(C)}) = \frac{1}{|\mathcal{C}(\mathcal{N})|} \sum_{C \in \mathcal{C}(\mathcal{N})} H_S^n(p^{v, C}) \\ &= H_{HG}^{n, v}(v). \end{aligned}$$

(Sufficiency). The proof goes as follows. In part (i), we show that (HG) holds for any totally ordered regular set system  $C$ , i.e., we show that  $H^{n, C} = H_{HG}^{n, C}$  for any  $n \geq 2$ . In part (ii), we extend the result to any regular set system, i.e.,  $H^{n, v} = H_{HG}^{n, v}$  on  $\mathcal{A}$ .

(i) We show by induction on  $n$  that for any  $v$  on a totally ordered regular set system  $(N, C)$ ,  $C$  being a maximal chain:

$$H^{n, C}(v) = H_{HG}^{n, C} = H_S^n(p^{v, C}). \tag{5.1}$$

First, we show the result for  $n = 2$ , using (HG1), (HG2) and (HG3), in a way similar to the Faddeev's axiomatization for the Shannon entropy. Suppose that  $v^3$  is a capacity in  $(N := \{1, 2, 3\}, \mathbf{3} := \{\emptyset, \{1\}, \{1, 2\}, N\})$  and  $v^3(1) := s$ ,  $v^3(\{1, 2\}) := s + t$ ,  $v^3(N) := s + t + u = 1$ ,  $s, u \geq 0$ ,  $t > 0$ . We can regard

$v^3$  as the embedding of  $v^2 = (0, t/(1-s), 1)$  into  $v = (0, s, 1)$  on  $\mathcal{N} := \{\emptyset, \{1\}, \{1, 2\}\}$  at  $\{1, 2\}$  (putting  $i'_k = 2, i''_k = 3$  with notations of Definition 8). Then, by (HG3), we get

$$\begin{aligned} H^{3,3}(v^3) &= H^{2,v}(0, s, 1) + (1-s)H^{2,2}(0, t/(1-s), 1) \\ &= H^{2,v}(0, s, 1) + (1-s)H^{2,2}(0, (1-s-t)/(1-s), 1) \\ &= H^{2,v}(0, s, 1) + (1-s)H^{2,2}(0, u/(1-s), 1), \end{aligned} \tag{5.2}$$

where the second equality follows from (HG2) and  $s+t+u=1$ . Similarly, we can also regard  $v^3$  as the embedding of  $v^2 = (0, s/(s+t), 1)$  into  $v = (0, s+t, 1)$  on  $\mathcal{N}' := \{\emptyset, \{1\}, \{1, 3\}\}$  at  $\{1\}$  (putting  $i'_k = 1, i''_k = 2$ ). This time, we obtain by (HG3) and  $s+t+u=1$ ,

$$\begin{aligned} H^{3,3}(v^3) &= H^{2,v'}(0, s+t, 1) + (s+t)H^{2,2}(0, s/(s+t), 1) \\ &= H^{2,v'}(0, u, 1) + (1-u)H^{2,2}(0, s/(1-u), 1). \end{aligned} \tag{5.3}$$

Hence (5.2) and (5.3) yield

$$H^{2,v}(0, s, 1) + (1-s)H^{2,2}(0, u/(1-s), 1) = H^{2,v'}(0, u, 1) + (1-u)H^{2,2}(0, s/(1-u), 1).$$

Since regular set systems  $\mathcal{N}, \mathcal{N}'$  and  $\mathbf{2}$  are isomorphic, putting  $f(s) := H^{2,2}(0, s, 1)$ , we obtain

$$f(s) + (1-s)f(u/(1-s)) = f(u) + (1-u)f(s/(1-u)), \tag{5.4}$$

where the above expression is valid for any  $s, u \in [0, 1)$  satisfying  $s+u < 1$ . If  $s=0, u > 0$ , then

$$f(0) + f(u) = f(u) + (1-u)f(0),$$

hence we obtain

$$f(0) = H^{2,2}(0, 0, 1) = H^{2,2}(0, 1, 1) = 0.$$

Integrating both sides of (5.4) for  $u$  from 0 to  $1-s$ , where  $0 \leq s < 1$ , we obtain

$$(1-s)f(s) + (1-s)^2 \int_0^1 f(x)dx = \int_0^{1-s} f(x)dx + s^2 \int_s^1 x^{-3}f(x)dx. \tag{5.5}$$

By (HG1),  $f(s)$  is continuous on  $[0, 1]$ , hence all terms except the first one of the left side of (5.5) are differentiable on  $0 < s < 1$ , so that  $f(s)$  is also differentiable on  $0 < s < 1$ . Therefore differentiating (5.5) with respect to  $s$ , we have

$$(1-s)f'(s) - f(s) - 2(1-s) \int_0^1 f(x)dx = -f(1-s) + 2s \int_s^1 x^{-3}f(x)dx - \frac{f(s)}{s}.$$

Since  $f(s) = f(1-s)$ ,

$$(1-s)f'(s) = 2(1-s) \int_0^1 f(x)dx + 2s \int_s^1 x^{-3}f(x)dx - \frac{f(s)}{s}. \tag{5.6}$$

By the same arguments as above,  $f'(s)$  is differentiable on  $0 < s < 1$ , then differentiating (5.6) by  $s$ , we have for  $0 < s < 1$ ,

$$f''(s) = \frac{-2}{s(1-s)} \int_0^1 f(x)dx. \tag{5.7}$$

Integrating (5.7) twice, we obtain for  $0 \leq s \leq 1$ ,

$$f(s) = -s \log s - (1-s) \log(1-s)$$

by  $f(0) = 0$  and  $f(1/2) = H^{2,2}(0, 1/2, 1) = 1$ . Therefore, for any  $v^2$  on  $\mathbf{2}$ ,

$$H^{2,2}(v^2) = H^2_S(p^{v^2,C}) = H^2_{HG}(v^2).$$



Suppose that (5.1) holds till  $n$  and show that it still holds for  $n + 1$ . Let  $v$  be a capacity on a maximal chain  $C := (c_0, \dots, c_n)$ . Then by (HG3), we have, denoting by  $C'$  the chain after embedding:

$$\begin{aligned} H^{n+1,C'}(v^{c_k}) &= H^{n,C}(v) + (v(c_k) - v(c_{k-1}))H^{2,2}(v^2) = H_S^n(p^{v,C}) + (v(c_k) - v(c_{k-1}))H^{2,2}(0, u, 1) \\ &= H_S^n(p^{v,C}) + (v(c_k) - v(c_{k-1}))H^{2,2}\left(0, \frac{v^{c_k}(c'_k) - v^{c_k}(c_{k-1})}{v(c_k) - v(c_{k-1})}, 1\right) \\ &= H_S^n(p^{v,C}) + (v(c_k) - v(c_{k-1}))H_S^2\left(\frac{v^{c_k}(c'_k) - v^{c_k}(c_{k-1})}{v(c_k) - v(c_{k-1})}, \frac{v^{c_k}(c'_k) - v^{c_k}(c'_k)}{v(c_k) - v(c_{k-1})}\right) = H_S^{n+1}(p^{v^{c_k},C'}) \\ &= H_{HG}^{n+1,C'}(v^{c_k}) \end{aligned}$$

by (3.1) and (F3), which means (5.1) holds for  $v^{c_k}$ , so that (HG) holds for any capacities on totally ordered regular set systems. From now on, the expression of the entropy on totally ordered regular set systems of size  $n$  will be denoted by  $H^{n,n}$ .

(ii) We show that for a regular set system which is not totally ordered, (HG) holds.  $2^N$  has  $n!$  maximal chains  $C_1, \dots, C_{n!}$ . By (HG4), there exist unique  $\alpha_{C_1}, \dots, \alpha_{C_{n!}} > 0$ ,  $\alpha_{C_1} + \dots + \alpha_{C_{n!}} = 1$ , such that for any capacity  $v$  on  $2^N$  it holds

$$H^{n,2^N}(v) = \alpha_{C_1}H^{n,n}(v|_{C_1}) + \dots + \alpha_{C_{n!}}H^{n,n}(v|_{C_{n!}}). \tag{5.8}$$

Let  $u_1, \dots, u_n > 0$ , and put  $h_S(u) := -u \log u - (1 - u) \log(1 - u)$ . First we define a capacity  $v$  on  $2^N$  by

$$v(A) := \begin{cases} 0, & |A| < 1, \\ u_i, & A = \{i\}, \quad i = 1, \dots, n, \\ 1, & |A| > 1. \end{cases}$$

Then,

$$H^{n,2^N}(v) = \left(\sum_{C \ni \{1\}} \alpha_C\right) h_S(u_1) + \dots + \left(\sum_{C \ni \{n\}} \alpha_C\right) h_S(u_n).$$

For any permutation  $\pi$  on  $N$  and any chain  $C_i$ , we have

$$\pi \circ v(A) = \begin{cases} 0, & A = \emptyset \\ u_{\pi^{-1}(i)}, & A = \{i\}, \quad i = 1, \dots, n, \\ 1, & |A| > 1, \end{cases}$$

hence

$$\begin{aligned} H^{n,2^N}(\pi \circ v) &= \alpha_{C_1}H^{n,n}((\pi \circ v)|_{C_1}) + \dots + \alpha_{C_{n!}}H^{n,n}((\pi \circ v)|_{C_{n!}}) \\ &= \left(\sum_{C \ni \{1\}} \alpha_C\right) h_S(u_{\pi^{-1}(1)}) + \dots + \left(\sum_{C \ni \{n\}} \alpha_C\right) h_S(u_{\pi^{-1}(n)}). \end{aligned}$$

By (HG5),  $H^{n,2^N}(v) = H^{n,2^N}(\pi \circ v)$  holds for any permutation  $\pi$  on  $N$ , and any  $u_1, \dots, u_n > 0$ , so that we obtain

$$\left(\sum_{C \ni \{1\}} \alpha_C\right) = \dots = \left(\sum_{C \ni \{n\}} \alpha_C\right) = \frac{1}{n}.$$

Next we consider the set system  $\mathcal{N}_1 := \{A \in 2^N \mid A \subsetneq \{1\}, A \supsetneq \{1\} \text{ or } A = \{1\}\}$ , which is a regular set system and has  $(n - 1)!$  maximal chains. Without loss of generality, we assume  $\mathcal{C}(\mathcal{N}_1) = \{C_1, \dots, C_{(n-1)!}\} \subsetneq \mathcal{C}(2^N)$ . Then by (HG4), there exist unique  $\beta_{C_1}, \dots, \beta_{C_{(n-1)!}} > 0$ ,  $\beta_{C_1} + \dots + \beta_{C_{(n-1)!}} = 1$ , such that for any capacity  $v$  on  $2^N$

$$H^{n,(n-1)!}(v|_{\mathcal{N}_1}) = \beta_{C_1}H^{n,n}(v|_{C_1}) + \dots + \beta_{C_{(n-1)!}}H^{n,n}(v|_{C_{(n-1)!}}). \tag{5.9}$$

Since  $\mathcal{C}(2^N) = \mathcal{C}(\mathcal{N}_1) \cup \{C_{(n-1)!+1}\} \cup \dots \cup \{C_{n!}\}$  and  $\mathcal{C}(\mathcal{N}_1), \{C_{(n-1)!+1}\}, \dots, \{C_{n!}\}$  are mutually disjoint, by applying (HG4) again, there exist  $\beta, \beta_{C_{(n-1)!+1}}, \dots, \beta_{C_{n!}} > 0$ ,  $\beta + \beta_{C_{(n-1)!+1}} + \dots + \beta_{C_{n!}} = 1$  such that for any capacity  $v$  on  $2^N$

$$H^{n,2^N}(v) = \beta H^{n,\mathcal{V}_1}(v|_{\mathcal{N}_1}) + \beta_{C_{(n-1)+1}} H^{n,n}(v|_{C_{(n-1)+1}}) + \dots + \beta_{C_n} H^{n,n}(v|_{C_n}).$$

Substituting (5.9) in the above and comparing with (5.8), we get in particular  $\alpha_{C_i} = \beta\beta_{C_i}$  for  $i = 1, \dots, (n-1)!$ . Since  $C_i \ni \{1\}, i = 1, \dots, (n-1)!$ ,

$$\sum_{i=1}^{(n-1)!} \beta\beta_{C_i} = \sum_{i=1}^{(n-1)!} \alpha_{C_i} = \sum_{C \ni \{1\}} \alpha_C = \frac{1}{n},$$

which implies  $\beta = 1/n$ . Let us take  $v$  on  $2^N$  such that

$$v|_{\mathcal{N}_1}(A) := \begin{cases} 0, & |A| < 2, \\ u_i, & A = \{1, i\}, i = 2, \dots, n, \\ 1, & |A| > 2. \end{cases}$$

Then

$$H^{n,\mathcal{V}_1}(v|_{\mathcal{N}_1}) = \left( \sum_{C \ni \{1\}, \{1,2\}} \beta_C \right) h_S(u_2) + \dots + \left( \sum_{C \ni \{1\}, \{1,n\}} \beta_C \right) h_S(u_n),$$

and by (HG5),  $H^{n,\mathcal{V}_1}(v|_{\mathcal{N}_1}) = H^{n,\pi(\mathcal{V}_1)}(\pi \circ (v|_{\mathcal{N}_1}))$  holds for any permutation  $\pi$  on  $N$  satisfying  $\pi(\mathcal{N}_1) = \mathcal{N}_1$  (i.e., permutations on  $N \setminus \{1\}$ ), so that we obtain

$$\sum_{C \ni \{1\}, \{1,2\}} \beta_C = \dots = \sum_{C \ni \{1\}, \{1,n\}} \beta_C = \frac{1}{n-1}, \tag{5.10}$$

which implies for  $i = 2, \dots, n$ ,

$$\sum_{C \ni \{1\}, \{1,i\}} \alpha_C = \sum_{C \ni \{1\}, \{1,i\}} \beta\beta_C = \frac{1}{n(n-1)}.$$

Applying the same method for  $\mathcal{N}_i := \{A \in 2^X \mid A \subsetneq \{i\}, A \supsetneq \{i\} \text{ or } A = \{i\}\}, i = 2, \dots, n$ , and

$$v|_{\mathcal{N}_i}(A) := \begin{cases} 0, & |A| < 2, \\ u_j, & A = \{i, j\}, i \neq j, \\ 1, & |A| > 2, \end{cases}$$

we obtain for every  $i$  and  $j, i \neq j$ ,

$$\sum_{C \ni \{i\}, \{i,j\}} \alpha_C = \sum_{C \ni \{i\}, \{i,j\}} \beta\beta_C = \frac{1}{n(n-1)}.$$

Next, we consider the set system  $\mathcal{N}_{1,2} := \{A \in 2^X \mid A \subsetneq \{1\}, A \supsetneq \{1,2\} \text{ or } A = \{1\}, \{1,2\}\}$ , and we assume w.l.o.g. that  $\mathcal{C}(\mathcal{N}_{1,2}) := \{C_1, \dots, C_{(n-2)!}\}$ . Applying (HG4) twice, there exist unique positive  $\gamma_{C_1}, \dots, \gamma_{C_{(n-2)!}}, \gamma, \gamma_{C_{(n-2)!+1}}, \dots, \gamma_{C_{(n-1)!}}, \gamma_{C_1} + \dots + \gamma_{C_{(n-2)!}} = 1$ , and  $\gamma + \gamma_{C_{(n-2)!+1}} + \dots + \gamma_{C_{(n-1)!}} = 1$ , such that for any capacity  $v$  on  $2^N$  it holds

$$\begin{aligned} H^{n,\mathcal{V}_{1,2}}(v|_{\mathcal{N}_{1,2}}) &= \gamma_{C_1} H^{n,n}(v|_{C_1}) + \dots + \gamma_{C_{(n-2)!}} H^{n,n}(v|_{C_{(n-2)!}}) \\ H^{n,\mathcal{V}_1}(v|_{\mathcal{N}_1}) &= \gamma H^{n,\mathcal{V}_{1,2}}(v|_{\mathcal{N}_{1,2}}) + \gamma_{C_{(n-2)!+1}} H^{n,n}(v|_{C_{(n-2)!+1}}) + \dots + \gamma_{C_{(n-1)!}} H^{n,n}(v|_{C_{(n-1)!}}). \end{aligned}$$

Substituting the first equation into the second and comparing with (5.9), we get in particular  $\beta_{C_i} = \gamma\gamma_{C_i}$ , hence  $\alpha_{C_i} = \beta\gamma\gamma_{C_i}$ , for  $i = 1, \dots, (n-2)!$ . Since  $C_i \ni \{1\}, \{1,2\}, i = 1, \dots, (n-2)!$ ,

$$\sum_{i=1}^{(n-2)!} \beta\gamma\gamma_{C_i} = \sum_{i=1}^{(n-2)!} \beta\beta_{C_i} = \sum_{C \ni \{1\}, \{1,2\}} \beta\beta_C = \frac{1}{n(n-1)},$$

which implies  $\gamma = 1/(n-1)$ . Taking a capacity  $v$  on  $2^N$  such that

$$v|_{\{1,2\}}(A) := \begin{cases} 0, & |A| < 3, \\ u_i, & A = \{1, 2, i\}, \quad i = 3, \dots, n, \\ 1, & |A| > 3, \end{cases}$$

we have

$$H^{n-1,2}(v|_{\{1,2\}}) = \left( \sum_{C \ni \{1\}, \{1,2\}, \{1,2,3\}} \gamma_C \right) h_S(u_3) + \dots + \left( \sum_{C \ni \{1\}, \{1,2\}, \{1,2,n\}} \gamma_C \right) h_S(u_n).$$

By (HG5), using  $\pi$  on  $N$  satisfying  $\pi(\mathcal{N}_{1,2}) = \mathcal{N}_{1,2}$  (i.e., the permutations on  $N \setminus \{1, 2\}$ ), we obtain

$$\sum_{C \ni \{1\}, \{1,2\}, \{1,2,3\}} \gamma_C = \dots = \sum_{C \ni \{1\}, \{1,2\}, \{1,2,n\}} \gamma_C = \frac{1}{n-2},$$

which implies for  $i = 3, \dots, n$ , and using  $\beta = 1/n$  and  $\gamma = 1/(n-1)$ , we have

$$\sum_{C \ni \{1\}, \{1,2\}, \{1,2,i\}} \alpha_C = \sum_{C \ni \{1\}, \{1,2\}, \{1,2,i\}} \beta \gamma \gamma_C = \frac{1}{n(n-1)(n-2)}.$$

Applying the same method for  $\mathcal{N}_{i,j} := \{A \in 2^N | A \subsetneq \{i\}, A \supseteq \{i, j\} \text{ or } A = \{i, \{i, j\}\}, i \neq j\}$ , we obtain for every  $i, j, k$  satisfying  $i \neq j, i \neq k, j \neq k$ ,

$$\sum_{C \ni \{i\}, \{i,j\}, \{i,j,k\}} \alpha_C = \sum_{C \ni \{i\}, \{i,j\}, \{i,j,k\}} \beta \gamma \gamma_C = \frac{1}{n(n-1)(n-2)}.$$

Repeating a similar process up to  $\mathcal{N}_{i_1, \dots, i_{n-1}} := \{A \in 2^N | A \subsetneq \{i_1\}, A \supseteq \{i_1, \dots, i_{n-1}\} \text{ or } A = \{i_1, \{i_1, i_2\}, \dots, \{i_1, \dots, i_{n-1}\}\}$ , we finally obtain  $\alpha_{C_i} = 1/n!, i = 1, \dots, n!$ , that is,

$$H^{n,2^N}(v) = \frac{1}{n!} \sum_{C \in \mathcal{C}(2^N)} H^{n,n}(v|_C).$$

Consider a regular set system  $\mathcal{N} \subsetneq 2^N$ . We can write  $2^N$  as the union of all its  $n!$  maximal chains:

$$2^N = C_1 \cup C_2 \cup \dots \cup C_{n!}.$$

For commodity, let us call  $C_1, \dots, C_k$  the chains which are not in  $\mathcal{N}$ , hence

$$\mathcal{C}(2^N) = \{C_1\} \cup \dots \cup \{C_k\} \cup \mathcal{C}(\mathcal{N}).$$

and  $\{C_1\}, \dots, \{C_k\}$  and  $\mathcal{C}(\mathcal{N})$  are mutually disjoint. By (HG4), we can find unique positive  $\alpha_1, \dots, \alpha_k, \alpha_{k+1}$  with  $\sum_{i=1}^{k+1} \alpha_i = 1$ , such that

$$H^{n,2^N}(v) = \alpha_1 H^{n,1}(v|_{C_1}) + \dots + \alpha_k H^{n,1}(v|_{C_k}) + \alpha_{k+1} H^{n,1}(v|_{\mathcal{N}}).$$

Similarly, there exists unique positive  $\beta_{k+1}, \dots, \beta_{n!}$  with  $\sum_{i=k+1}^{n!} \beta_i = 1$ , such that

$$H^{n,1}(v|_{\mathcal{N}}) = \beta_{k+1} H^{n,1}(v|_{C_{k+1}}) + \dots + \beta_{n!} H^{n,1}(v|_{C_{n!}}).$$

Substituting in the above leads to

$$H^{n,2^N}(v) = \alpha'_1 H^{n,1}(v|_{C_1}) + \dots + \alpha'_k H^{n,1}(v|_{C_k}) + \alpha'_{k+1} H^{n,1}(v|_{C_{k+1}}) + \dots + \alpha'_{n!} H^{n,1}(v|_{C_{n!}}),$$

with  $\alpha'_i = \alpha_i$  for  $i = 1, \dots, k$ , and  $\alpha'_i = \alpha_{k+1} \beta_i$  for  $i = k+1, \dots, n!$ . We have proved that  $\alpha'_1 = \alpha'_2 = \dots = \alpha'_{n!} = 1/n!$ , for  $i = 1, \dots, n!$ . It follows that  $k/n! + \alpha_{k+1} = 1$ , hence

$$\alpha_{k+1} = \frac{n! - k}{n!} = \frac{|\mathcal{C}(\mathcal{N})|}{n!}$$

by definition of  $C_{k+1}, \dots, C_{n!}$ . Now,  $\alpha_{k+1} \beta_i = 1/n!$  for  $i = k+1, \dots, n!$ , hence  $\beta_i = 1/|\mathcal{C}(\mathcal{N})|$ , for  $i = k+1, \dots, n!$ . Finally, we get

$$H^{n,\mathcal{N}}(v|_{\mathcal{N}'} ) = \frac{1}{|\mathcal{C}(\mathcal{N}')|} \sum_{i=k+1}^{n!} H^{n,1}(v|_{C_i}) = \frac{1}{|\mathcal{C}(\mathcal{N}')|} \sum_{C \in \mathcal{C}(\mathcal{N}')} H_S(p^{v,C})$$

using (5.1).  $\square$

**Proof of Theorem 16 (Necessity).** When  $v$  is a capacity on a totally ordered regular set system,  $H_D(v) = H_{HG}(v)$  holds. Hence (D) implies (HG1), (HG2) and (HG3) by Theorem 14. We show that (D) satisfies (HG4'). We have with set systems  $\mathcal{N}_1, \dots, \mathcal{N}_k$  satisfying conditions of (HG4'),

$$\begin{aligned} H_D^{n,\mathcal{N}}(v) &= \min_{C \in \mathcal{C}(\mathcal{N}')} H_S^n(p^{v,C}) = \min \left\{ \min_{C \in \mathcal{C}(\mathcal{N}_1)} H_S^n(p^{v,C}), \dots, \min_{C \in \mathcal{C}(\mathcal{N}_k)} H_S^n(p^{v,C}) \right\} \\ &= \min \{ H_D^{n,\mathcal{N}_1}(v_1), \dots, H_D^{n,\mathcal{N}_k}(v_k) \} \end{aligned}$$

with usual notations.

(Sufficiency). When  $v$  is a capacity on a totally ordered regular set system,  $H_D(v) = H_{HG}(v)$  holds. Hence (HG1), (HG2) and (HG3) imply (D) by the same way of the Proof of Theorem 14.

Next, let  $v$  be a capacity on  $\mathcal{N}$  which is regular and not totally ordered, and write  $\mathcal{N}$  as the union of all its maximal chains:  $\mathcal{N} = C_1 \cup \dots \cup C_k$ . Then by application of (HG4') we have

$$\begin{aligned} H^{n,\mathcal{N}}(v) &= \min \{ H^{n,C_1}(v|_{C_1}), \dots, H^{n,C_k}(v|_{C_k}) \} = \min \{ H_S^n(p^{v|_{C_1},C_1}), \dots, H_S^n(p^{v|_{C_k},C_k}) \} = \min_{C \in \mathcal{C}(\mathcal{N}')} H_S^n(p^{v,C}), \\ &= H_D^{n,\mathcal{N}}(v). \quad \square \end{aligned}$$

### 6. Applications and concluding remarks

We have proposed an axiomatization of the entropy of capacities defined on regular set systems, in a spirit very close to the axiomatization of Faddeev for the Shannon entropy. The key axiom which is additional is (HG4), binding the computation of entropy to the structure of the set system.

Essentially, two comments are of interest here. The first one concerns the nature of the five axioms we propose. Axioms (HG1), (HG3) and (HG5) clearly correspond to the three axioms of Faddeev for the Shannon entropy. Axiom (HG2) is necessary in our framework, but is implicit for the Shannon entropy. Put differently, (HG2) too is characteristic of Shannon entropy. Hence, only axiom (HG4) is specific to our construction; it is able to deal with the structure of the set system, provided it is regular. It is responsible for the general form of the entropy, that is, a weighted average of classical Shannon entropies along maximal chains. This suggests the following: as other entropies have been proposed for probabilities, together with axiomatizations, it would be possible to carry the present framework for axiomatizing these other entropies, just replacing axioms (HG1), (HG2), (HG3) and (HG5) by axioms which are specific to these entropies.

Another comment concerns the usefulness of regular set systems. Would it be possible to consider yet more general structures, i.e., set systems which are not regular? Formally, Definition 11 can be applied to any set system, since maximal chains are always defined, and thus probability distributions  $p^{v,C}$  too. A fundamental problem is however that these chains need not to be of the same length, whose direct consequence is that the probability distribution would be undefined for some elements of  $N$ . For example, take  $\mathcal{N}' := \{\emptyset, \{1\}, \{1, 2, 3\}\}$ , which is totally ordered but not regular, and let  $v$  be a capacity defined on  $\mathcal{N}'$ . Computing the entropy of  $v$  by Definition 11 leads to

$$H(v) = -v(\{1\}) \log v(\{1\}) - (v(\{1, 2, 3\}) - v(\{1\})) \log (v(\{1, 2, 3\}) - v(\{1\})).$$

Clearly, the quantities  $v(\{1\})$ ,  $v(\{1, 2, 3\}) - v(\{1\})$  cannot be considered as the values of a probability distribution on  $N$ . Another troublesome fact is the following. We know that classically the entropy takes its maximal value when  $v$  is an additive uniform capacity, that is, which satisfies  $v(A) = |A|/n$ . But for  $v$  on  $\mathcal{N}'$ ,  $H(v)$  takes the maximal value when  $v(\{1\}) = 1/2$  which is not the additive uniform capacity.

It seems then, as far as maximal chains are considered as fundamental ingredients, that regular set systems are the most general structure we can consider. It is important to note that regular set systems encompass

many well-known ordered structures. This is studied in detail in [8], and it turns out that distributive lattices, set lattices, convex geometries and antimatroids are strictly included into regular set systems. The following simple example shows that a regular set system is not necessarily a lattice: take  $N = \{1, 2, 3, 4\}$  and  $\mathcal{A} := \{\emptyset, \{1\}, \{3\}, \{1, 2\}, \{2, 3\}, \{1, 4\}, \{3, 4\}, \{1, 2, 3\}, \{1, 3, 4\}, N\}$ . Then,  $\{1\}$  and  $\{3\}$  have no supremum. Now, the case of lattices is obviously a particular case of interest. We refer the reader to [6] where we have examined the application of our definition of entropy to the case where capacities are defined on lattices (not necessarily set lattices).

### Acknowledgements

We address all our deepest thanks to the anonymous reviewers for a careful reading of the manuscript and the detection of several flaws in the previous version of the axiomatization.

### References

- [1] G. Choquet, Theory of capacities, *Annales de l'Institut Fourier* 5 (1953) 131–295.
- [2] D. Denneberg, *Non-additive Measure and Integral*, Academic, Kluwer, 1994.
- [3] A. Dukhovny, General entropy of general measures, *International Journal of Uncertainty Fuzziness Knowledge-Based Systems* 10 (2002) 213–225.
- [4] B.R. Ebanks, P.K. Sahoo, W. Sander, *Characterizations of Information Measures*, World Scientific, 1998.
- [5] D.K. Faddeev, The notion of entropy of finite probabilistic schemes (Russian), *Uspekhi Mat. Nauk* 11 (1956) 15–19.
- [6] A. Honda, M. Grabisch, Entropy of capacities on lattices, *Information Sciences* 176 (2006) 3472–3489.
- [7] I. Kojadinovic, J.-L. Marichal, M. Roubens, An axiomatic approach to the definition of the entropy of a discrete Choquet capacity, *Information Sciences* 172 (2005) 131–153.
- [8] F. Lange, M. Grabisch, Values on regular games under Kirchhoff's laws, *SIAM Journal of Discrete Mathematics*, submitted for publication.
- [9] J.-L. Marichal, Entropy of discrete Choquet capacities, *European Journal of Operation Research* 137 (2002) 612–624.
- [10] J.-L. Marichal, M. Roubens, Entropy of discrete fuzzy measure, *International Journal of Uncertainty Fuzziness Knowledge-Based Systems* 8 (2000) 625–640.
- [11] C.E. Shannon, A mathematical theory of communication, *Bell System Technical Journal* 27 (1948) 374–423, 623–656.
- [12] M. Sugeno, *Theory of Fuzzy Integrals and its Applications*. Doctoral Thesis, Tokyo Institute of Technology, 1974.
- [13] R.R. Yager, On the entropy of fuzzy measures, *IEEE Transaction on Fuzzy Systems* 8 (2000) 453–461.